



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/505	A1	(11) International Publication Number: WO 96/16657 (43) International Publication Date: 6 June 1996 (06.06.96)
(21) International Application Number: PCT/EP95/04065 (22) International Filing Date: 16 October 1995 (16.10.95) (30) Priority Data: 9423911.8 26 November 1994 (26.11.94) GB (71) Applicant (for GB only): PFIZER LIMITED [GB/GB]; Ramsgate Road, Sandwich, Kent CT13 9NJ (GB). (71) Applicant (for all designated States except GB JP US): PFIZER RESEARCH AND DEVELOPMENT COMPANY, N.V./S.A. [BE/IE]; La Touche House, International Financial Services Centre, Dublin 1 (IE). (71) Applicant (for JP only): PFIZER INC. [US/US]; 235 East 42nd Street, New York, NY 10017 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): CAMPBELL, Simon, Fraser [GB/GB]; Pfizer Central Research, Ramsgate Road, Sandwich, Kent CT13 9NJ (GB). (74) Agents: MOORE, James, William et al.; Pfizer Limited, European Patent Dept., Ramsgate Road, Sandwich, Kent CT13 9NJ (GB).		(81) Designated States: CA, JP, MX, US, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: BICYCLIC HETEROCYCLIC COMPOUNDS FOR THE TREATMENT OF IMPOTENCE (57) Abstract The use of certain 5-arylpyrazolo[4,3-d]pyrimidin-7-ones, 6-arylpyrazolo[3,4-d]pyrimidin-4-ones, 2-arylquinazolin-4-ones, 2-arylpyrido[3,2-d]pyrimidin-4-ones, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition containing either entity, for the manufacture of a medicament for the curative or prophylactic treatment of erectile dysfunction in a male animal, including man; a pharmaceutical composition for said treatment; and a method of said treatment of said animal with said pharmaceutical composition or with said either entity.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

BICYCLIC HETEROCYCLIC COMPOUNDS FOR
THE TREATMENT OF IMPOTENCE

This invention relates to the use of certain pyrazolo[4,3-d]pyrimidin-7-ones, pyrazolo[3,4-d]pyrimidin-4-ones, quinazolin-4-ones, purin-6-ones and pyrido[3,2-d]pyrimidin-4-ones for the treatment of impotence.

Impotence can be defined literally as a lack of power, in the male, to copulate and may involve an inability to achieve penile erection or ejaculation, or both. More specifically, erectile impotence or dysfunction may be defined as an inability to obtain or sustain an erection adequate for intercourse. Its prevalence is claimed to be between 2 and 7% of the human male population, increasing with age, up to 50 years, and between 18 and 75% between 55 and 80 years of age. In the USA alone, for example, it has been estimated that there are up to 10 million impotent males, with the majority suffering from problems of organic rather than of psychogenic origin.

Reports of well-controlled clinical trials in man are few and the efficacy of orally administered drugs is low. Although many different drugs have been shown to induce penile erection, they are only effective after direct injection into the penis, e.g. intraurethrally or intracavernosally (i.c.), and are not approved for erectile dysfunction. Current medical treatment is based on the i.c. injection of vasoactive substances and good results have been claimed with phenoxybenzamine, phentolamine, papaverine and prostaglandin E₁, either alone or in combination; however, pain, priapism and fibrosis of the penis are associated with the i.c. administration of some of these agents. Potassium channel openers (KCO) and vasoactive intestinal polypeptide (VIP) have also been shown to be active i.c., but cost and stability issues could limit development of the latter. An alternative


to the i.c. route is the use of glyceryl trinitrate (GTN) patches applied to the penis, which has been shown to be effective but produces side-effects in both patient and partner.

As a general alternative to pharmacological intervention, a variety of penile prostheses has been used to assist achievement of an erection. The short term success rate is good, but problems with infection and ischaemia, especially in diabetic men, make this type of treatment a final option rather than first-line therapy.

The compounds of the invention are potent inhibitors of cyclic guanosine 3',5'-monophosphate phosphodiesterases (cGMP PDEs) in contrast to their inhibition of cyclic adenosine 3',5'-monophosphate phosphodiesterases (cAMP PDEs). This selective enzyme inhibition leads to elevated cGMP levels which, in turn, provides the basis for the utilities already disclosed for the said compounds in WO-A-93/06104, WO-A-93/07149, WO-A-93/12095, WO-A-94/00453 and WO-A-94/05661 respectively, namely in the treatment of stable, unstable and variant (Prinzmetal) angina, hypertension, pulmonary hypertension, congestive heart failure, atherosclerosis, conditions of reduced blood vessel patency e.g. post-percutaneous transluminal coronary angioplasty (post-PTCA), peripheral vascular disease, stroke, bronchitis, allergic asthma, chronic asthma, allergic rhinitis, glaucoma, and diseases characterised by disorders of gut motility, e.g. irritable bowel syndrome (IBS).


Unexpectedly, it has now been found that these disclosed compounds are useful in the treatment of erectile dysfunction. Furthermore the compounds may be administered orally, thereby obviating the disadvantages associated with i.c. administration. Thus the present invention concerns the use of a compound of formula (I):

3



wherein R^1 is methyl or ethyl;
 R^2 is ethyl or n-propyl;
 and R^3 and R^4 are each independently H, or C_1-C_6
 alkyl optionally substituted with C_3-C_7
 cycloalkyl or with morpholino;

a compound of formula (II):




wherein R^1 is C_1-C_6 alkyl;
 R^2 is H; methyl or ethyl;
 R^3 is C_2-C_4 alkyl;
 R^4 is H; C_1-C_4 alkyl optionally substituted
 with NR^5R^6 , CN, $CONR^5R^6$ or CO_2R^7 ; C_2-C_4 alkenyl
 optionally substituted with CN, $CONR^5R^6$ or
 CO_2R^7 ; C_2-C_4 alkanoyl optionally substituted
 with NR^5R^6 ; $SO_2NR^5R^6$; $CONR^5R^6$; CO_2R^7 or halo;
 R^5 and R^6 are each independently H or C_1-C_4

alkyl; or, together with the nitrogen atom to which they are attached, form a pyrrolidino, piperidino, morpholino, 4-(NR⁸)-1-piperazinyl or 1-imidazolyl group wherein said group is optionally substituted with one or two C₁-C₄ alkyl groups;

R⁷ is H or C₁-C₄ alkyl;

and R⁸ is H; C₁-C₃ alkyl or (hydroxy)C₂-C₃ alkyl;

a compound of formula (III):



wherein R¹ is H; C₁-C₄ alkyl; C₁-C₄ alkoxy or CONR⁵R⁶;
 R² is H or C₁-C₄ alkyl;
 R³ is C₂-C₄ alkyl;
 R⁴ is H; C₂-C₄ alkanoyl optionally substituted with NR⁷R⁸; (hydroxy)C₂-C₄ alkyl optionally substituted with NR⁷R⁸; CH=CHCO₂R⁹; CH=CHCONR⁷R⁸; CH₂CH₂CO₂R⁹; CH₂CH₂CONR⁷R⁸; SO₂NR⁷R⁸; SO₂NH(CH₂)_nNR⁷R⁸ or imidazolyl;
 R⁵ and R⁶ are each independently H or C₁-C₄ alkyl;
 R⁷ and R⁸ are each independently H or C₁-C₄ alkyl; or, together with the nitrogen atom to which they are attached, form a pyrrolidino, piperidino, morpholino or 4-(NR¹⁰)-1-piperazinyl group wherein any

5

of said groups is optionally substituted with CONR^5R^6 ;


R^9 is H or $\text{C}_1\text{-C}_4$ alkyl;

R^{10} is H; $\text{C}_1\text{-C}_3$ alkyl or (hydroxy) $\text{C}_2\text{-C}_3$ alkyl;

and n is 2, 3 or 4;

with the proviso that R^4 is not H when R^1 is H, $\text{C}_1\text{-C}_4$ alkyl or $\text{C}_1\text{-C}_4$ alkoxy;

a compound of formula (IV):



wherein R^1 is $\text{C}_1\text{-C}_4$ alkyl;


R^2 is $\text{C}_2\text{-C}_4$ alkyl;

R^3 is H or $\text{SO}_2\text{NR}^4\text{R}^5$;

R^4 and R^5 together with the nitrogen atom to which they are attached form a pyrrolidino, piperidino, morpholino or 4-(NR^6)-1-piperazinyl group;

and R^6 is H or $\text{C}_1\text{-C}_3$ alkyl;

or a compound of formula (V):



wherein R^1 is H; C_1 - C_4 alkyl; CN or $CONR^4R^5$;
 R^2 is C_2 - C_4 alkyl;
 R^3 is $SO_2NR^6R^7$; NO_2 ; NH_2 ; $NHCOR^8$; $NHSO_2R^8$ or $N(SO_2R^8)_2$;
 R^4 and R^5 are each independently selected from H and C_1 - C_4 alkyl;
 R^6 and R^7 are each independently selected from H and C_1 - C_4 alkyl optionally substituted with CO_2R^9 , OH, pyridyl, 5-isoxazolin-3-onyl, morpholino or 1-imidazolidin-2-onyl; or, together with the nitrogen atom to which they are attached, form a pyrrolidino, piperidino, morpholino, 1-pyrazolyl or 4-(NR^{10})-1-piperazinyl group wherein any of said groups may optionally be substituted with one or two substituents selected from C_1 - C_4 alkyl, CO_2R^9 , NH_2 and OH;
 R^8 is C_1 - C_4 alkyl or pyridyl;
 R^9 is H or C_1 - C_4 alkyl;
and R^{10} is H; C_1 - C_4 alkyl or (hydroxy) C_2 - C_3 alkyl;

or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition containing either entity, for the manufacture of a medicament for the curative or prophylactic treatment of erectile dysfunction in a

male animal, including man.

In the above definition, unless otherwise indicated, alkyl and alkoxy groups having three or more carbon atoms, and alkenyl and alkanoyl groups having four carbon atoms, may be straight chain or branched chain. Halo means fluoro, chloro, bromo or iodo.

The compounds of the invention may contain one or more asymmetric centres and thus they can exist as enantiomers or diastereoisomers. Furthermore, certain compounds of formulae (II) and (III) which contain alkenyl groups may exist as cis-isomers or trans-isomers. In each instance, the invention includes both mixtures and separate individual isomers.

The compounds of the invention may also exist in tautomeric forms and the invention includes both mixtures and separate individual tautomers.

The pharmaceutically acceptable salts of the compounds of the invention which contain a basic centre are, for example, non-toxic acid addition salts formed with inorganic acids such as hydrochloric, hydrobromic, sulphuric and phosphoric acid, with organo-carboxylic acids, or with organo-sulphonic acids. The compounds of the invention can also provide pharmaceutically acceptable metal salts, in particular non-toxic alkali metal salts, with bases. Examples include the sodium and potassium salts. For a review on suitable pharmaceutical salts, see J. Pharm. Sci., 1977, 66, 1.

A preferred group of compounds is that of formula (I) wherein R^3 is H; methyl or ethyl; R^4 is C_1 - C_6 alkyl optionally substituted with cyclohexyl or with morpholino; and R^1 and R^2 are as previously defined for formula (I); of formula (II) wherein R^1 is n-propyl; R^2 is H or methyl; R^3 is ethyl or n-propyl; R^4 is H; ethyl substituted with $CONR^5R^6$ or CO_2R^7 ; vinyl substituted with $CONR^5R^6$ or CO_2R^7 ; acetyl substituted with NR^5R^6 ; $SO_2NR^5R^6$; $CONR^5R^6$; CO_2R^7 or bromo; R^5 and R^6

together with the nitrogen atom to which they are attached form a morpholino, 4-(NR⁸)-1-piperazinyl or 2,4-dimethyl-1-imidazolyl group; R⁷ is H or t-butyl; and R⁸ is methyl or 2-hydroxyethyl; of formula (III) wherein R¹ is H; methyl; methoxy or CONR⁵R⁶; R² is H or methyl; R³ is ethyl or n-propyl; R⁴ is H; acetyl optionally substituted with NR⁷R⁸; hydroxyethyl substituted with NR⁷R⁸; CH=CHCO₂R⁹; CH=CHCONR⁷R⁸; CH₂CH₂CO₂R⁹; SO₂NR⁷R⁸; SO₂NH(CH₂)₃NR⁷R⁸ or 1-imidazolyl; R⁵ and R⁶ are each independently H or ethyl; R⁷ and R⁸ together with the nitrogen atom to which they are attached form a piperidino, 4-carbamoylpiperidino, morpholino or 4-(NR¹⁰)-1-piperazinyl group; R⁹ is H or t-butyl; and R¹⁰ is H; methyl or 2-hydroxyethyl; with the proviso that R⁴ is not H when R¹ is H, methyl or methoxy; of formula (IV) wherein R¹ and R² are each independently ethyl or n-propyl; R⁴ and R⁵ together with the nitrogen atom to which they are attached form a 4-(NR⁶)-1-piperazinyl group; and R³ and R⁶ are as previously defined for formula (IV); and of formula (V) wherein R¹ is H; n-propyl; CN or CONH₂; R² is ethyl; R³ is SO₂NR⁶R⁷; NO₂; NH₂; NHCOCH(CH₃)₂; NHSO₂CH(CH₃)₂; NHSO₂(3-pyridyl) or N[SO₂(3-pyridyl)]₂; R⁶ is H; methyl or 2-hydroxyethyl; R⁷ is methyl optionally substituted with 2-pyridyl or 5-isoxazolin-3-onyl; or ethyl 2-substituted with OH, CO₂CH₂CH₃, morpholino or 1-imidazolidin-2-onyl; or R⁶ and R⁷ together with the nitrogen atom to which they are attached form a (4-CO₂R⁹)piperidino, 5-amino-3-hydroxy-1-pyrazolyl or 4-(NR¹⁰)-1-piperazinyl group; R⁹ is H or ethyl; and R¹⁰ is H; methyl or 2-hydroxyethyl.

A particularly preferred group of compounds is that of formula (III) wherein R¹ is methyl; CONH₂ or CONHCH₂CH₃; R² is H; R³ is ethyl or n-propyl; R⁴ is H; acetyl; 1-hydroxy-2-(NR⁷R⁸)ethyl; CH=CHCO₂C(CH₃)₃; CH=CHCONR⁷R⁸; SO₂NR⁷R⁸ or 1-imidazolyl; R⁷ and R⁸

together with the nitrogen atom to which they are attached form a 4-(NR¹⁰)-1-piperazinyl group; and R¹⁰ is methyl or 2-hydroxyethyl; with the proviso that R⁴ is not H when R¹ is methyl; of formula (IV) wherein R¹ is n-propyl; R² is ethyl; and R³ is 1-piperazinylsulphonyl or 4-methyl-1-piperazinylsulphonyl; and of formula (V) wherein R¹ is n-propyl or CN; R² is ethyl; R³ is SO₂NR⁶R⁷; NHSO₂CH(CH₃)₂; NHSO₂(3-pyridyl) or N[SO₂(3-pyridyl)]₂; R⁶ is H or methyl; R⁷ is methyl; or ethyl 2-substituted with CO₂CH₂CH₃; morpholino or 1-imidazolidin-2-onyl; or R⁶ and R⁷ together with the nitrogen atom to which they are attached form a (4-CO₂R⁹)piperidino or 4-(NR¹⁰)-1-piperazinyl group; R⁹ is H or ethyl; and R¹⁰ is H; methyl or 2-hydroxyethyl.

Especially preferred individual compounds of the invention include:

1-ethyl-5-[5-(n-hexylsulphamoyl)-2-n-propoxyphenyl]-3-methyl-1,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one;

1-ethyl-5-(5-diethylsulphamoyl-2-n-propoxyphenyl)-3-methyl-1,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one;

5-[5-(N-cyclohexylmethyl-N-methylsulphamoyl)-2-n-propoxyphenyl]-1-ethyl-3-methyl-1,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one;

6-(5-bromo-2-n-propoxyphenyl)-3-methyl-1-n-propyl-1,5-dihydro-4H-pyrazolo[3,4-d]pyrimidin-4-one;

3-methyl-6-(5-morpholinolulphonyl-2-n-propoxyphenyl)-1-n-propyl-1,5-dihydro-4H-pyrazolo[3,4-d]pyrimidin-4-one;

6-[5-(2-carboxyvinyl)-2-n-propoxyphenyl]-3-methyl-1-n-propyl-1,5-dihydro-4H-pyrazolo[3,4-d]pyrimidin-4-one;

6-[5-(2-t-butoxycarbonylvinyl)-2-n-propoxyphenyl]-3-methyl-1-n-propyl-1,5-dihydro-4H-pyrazolo[3,4-d]pyrimidin-4-one;

3-methyl-6-[5-(2-morpholinocarbonylvinyl)-2-n-propoxyphenyl]-1-n-propyl-1,5-dihydro-4H-pyrazolo[3,4-d]pyrimidin-4-one;

3-methyl-6-[5-(2-morpholinocarbonylethyl)-2-n-propoxyphenyl]-1-n-propyl-1,5-dihydro-4H-pyrazolo[3,4-d]pyrimidin-4-one;

2-{2-ethoxy-5-[4-(2-hydroxyethyl)-1-piperazinylsulphonyl]phenyl}-8-methylquinazolin-4-(3H)-one;

2-{5-[4-(2-hydroxyethyl)-1-piperazinylsulphonyl]-2-n-propoxyphenyl}-8-methylquinazolin-4(3H)-one;

8-methyl-2-{5-[2-(4-methyl-1-piperazinylcarbonyl)ethenyl]-2-n-propoxyphenyl}quinazolin-4(3H)-one;

8-carbamoyl-2-{2-ethoxy-5-[4-(2-hydroxyethyl)-1-piperazinylsulphonyl]phenyl}quinazolin-4(3H)-one;

8-ethylcarbamoyl-2-(2-n-propoxyphenyl)quinazolin-4(3H)-one;

2-[2-ethoxy-5-(4-ethoxycarbonylpiperidinosulphonyl)phenyl]-8-n-propylpyrido[3,2-d]pyrimidin-4(3H)-one;

2-[5-(4-carboxypiperidinosulphonyl)-2-ethoxyphenyl]-8-n-propylpyrido[3,2-d]pyrimidin-4(3H)-one;

2-{2-ethoxy-5-[4-(2-hydroxyethyl)-1-piperazinylsulphonyl]phenyl}-8-n-propylpyrido[3,2-d]pyrimidin-4(3H)-one;

and 2-{2-ethoxy-5-[(bis-3-pyridylsulphonyl)amino]phenyl}-8-n-propylpyrido[3,2-d]pyrimidin-4(3H)-one.

The compounds of formulae (I), (II), (III), (IV) and (V) and their pharmaceutically acceptable salts, processes for the preparation thereof, in vitro test methods for determining the cGMP PDE and cAMP PDE inhibitory activities thereof, pharmaceutical compositions thereof and routes of administration for human use, are described in WO-A-93/06104, WO-A-93/07149, WO-A-93/12095, WO-A-94/00453 and WO-A-94/05661 respectively, which are incorporated herein by

reference.

A preliminary investigation was carried out with a view to isolating and characterising the cyclic nucleotide PDEs of human corpus cavernosum, relaxation of which leads to penile erection. Studies of substrate specificity, response to activators and inhibitor sensitivity, have demonstrated that human corpus cavernosum contains three distinct PDE enzymes.

Methods

Fresh frozen human penis was obtained from IIAM (Pennsylvania). Tissue was thawed at room temperature, the corpus cavernosum was dissected from the penis to yield approximately 2-4 g of tissue and the following isolation protocol was followed. Tissue was coarsely chopped in ice-cold isotonic buffer (35 ml) containing 250mM sucrose, 1mM EDTA, 0.5mM PMSF and 20mM HEPES, pH 7.2, and the mixture subjected to brief (1 min.) treatment with a Silversen mixer/emulsifier. Homogenates were prepared using homogeniser tubes with teflon pestles and soluble fraction was prepared by centrifugation at 100,000 x g for 60 min. at 4°C. 10 ml of high speed supernatant was applied to a Pharmacia Mono Q anion exchange column (1 ml bed volume) equilibrated with buffer containing 1mM EDTA, 0.5 mM PMSF and 20mM HEPES, pH 7.2 (chromatography buffer). The column was then washed with 5 bed volumes of chromatography buffer, after which PDEs were eluted using a continuous gradient of 0-500mM NaCl (total volume 35 ml) and 1 ml fractions collected.

Column fractions were assayed for PDE activity using 500nM cGMP or 500nM cAMP as substrate. cAMP PDE activity was also determined in the presence of 1 μ M unlabelled cGMP and the PDE activity of selected fractions was determined in the presence of 10mM CaCl₂ and 10 units/ml bovine brain calmodulin. Appropriate fractions were pooled and stored at 4°C during the

course of the study.

Inhibition studies were performed using a substrate concentration of 500nM throughout. All inhibitors were dissolved in DMSO and concentration-response curves were constructed over the range 3×10^{-10} to 1×10^{-4} M in half log increments. IC_{50} values were calculated using the sigmoidal curve fitting algorithm of biostat.

Results

Human corpus cavernosum soluble PDEs were separated into three distinct fractions of activity. The first, fraction I, (designated by order of elution) represents the major PDE present and is highly selective for cGMP as substrate. This fraction was found to be insensitive to stimulation by calcium/calmodulin and was classified as PDE_v . Fraction II hydrolyses cGMP and cAMP, with the latter activity being stimulated in the presence of cGMP, and is classified as PDE_{II} , whilst fraction III is cAMP selective and this activity is inhibited in the presence of cGMP, consistent with PDE_{III} activity.

In order to further characterise the PDE isoenzymes present in the tissue, studies were performed using a variety of inhibitors. Inhibitor studies with fractions I and II were performed using cGMP as substrate, whilst fraction III studies utilised cAMP. These studies confirmed that fraction I corresponds to PDE_v , whilst fraction III was clearly identified as PDE_{III} ; fraction II (PDE_{II}) was relatively insensitive to all the inhibitors tested.

In summary, the above investigation identified three PDE isoenzymes in human corpus cavernosum tissue. The predominant PDE is the cGMP-specific PDE_v , whilst cGMP-stimulated cAMP PDE_{II} and cGMP-inhibited cAMP PDE_{III}

are also present.

Certain compounds of the invention have been tested in vitro and found to be potent and selective inhibitors of the cGMP-specific PDE_v. Thus relaxation of the corpus cavernosum tissue and consequent penile erection is presumably mediated by elevation of cGMP levels in the said tissue, by virtue of the PDE inhibitory profile of the compounds of the invention.

Although the compounds of the invention are envisaged primarily for the treatment of erectile dysfunction or male sexual dysfunction, they may also be useful for the treatment of female sexual dysfunction, including orgasmic dysfunction related to clitoral disturbances, and of premature labour and dysmenorrhea.

Generally, in man, oral administration of the compounds of the invention is the preferred route, being the most convenient and avoiding the disadvantages associated with i.c. administration. A preferred dosing regimen for a typical man is 5 to 75 mg of compound three times daily. In circumstances where the recipient suffers from a swallowing disorder or from impairment of drug absorption after oral administration, the drug may be administered parenterally, e.g. sublingually or buccally.

For veterinary use, a compound of the invention or a non-toxic salt thereof is administered as a suitably acceptable formulation in accordance with normal veterinary practice and the veterinary surgeon will determine the dosing regimen and route of administration which will be most appropriate for a particular male animal.

Thus the invention includes a pharmaceutical composition for the curative or prophylactic treatment of erectile dysfunction in a male animal, including man, comprising a compound of formula (I), (II), (III), (IV) or (V), or a pharmaceutically acceptable salt

thereof, together with a pharmaceutically acceptable diluent or carrier.


There is further provided a process for the preparation of a pharmaceutical composition for the curative or prophylactic treatment of erectile dysfunction in a male animal, including man, comprising formulating a compound of formula (I), (II), (III), (IV) or (V), or a pharmaceutically acceptable salt thereof, with a pharmaceutically acceptable diluent or carrier.

The invention also provides a method of treating a male animal, including man, to cure or prevent erectile dysfunction which comprises treating said male animal with an effective amount of a compound of formula (I), (II), (III), (IV) or (V), or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition containing either entity.

15


CLAIMS

1. The use of a compound of formula (I):



wherein R^1 is methyl or ethyl;
 R^2 is ethyl or n-propyl;
 and R^3 and R^4 are each independently H, or C_1-C_6 alkyl optionally substituted with C_5-C_7 cycloalkyl or with morpholino;

a compound of formula (II):




wherein R^1 is C_1-C_6 alkyl;
 R^2 is H; methyl or ethyl;
 R^3 is C_2-C_4 alkyl;
 R^4 is H; C_1-C_4 alkyl optionally substituted with NR^5R^6 , CN, $CONR^5R^6$ or CO_2R^7 ; C_2-C_4 alkenyl

optionally substituted with CN, CONR^5R^6 or CO_2R^7 ; $\text{C}_2\text{-C}_4$ alkanoyl optionally substituted with NR^5R^6 ; $\text{SO}_2\text{NR}^5\text{R}^6$; CONR^5R^6 ; CO_2R^7 or halo; R^5 and R^6 are each independently H or $\text{C}_1\text{-C}_4$ alkyl; or, together with the nitrogen atom to which they are attached, form a pyrrolidino, piperidino, morpholino, 4-(NR^8)-1-piperazinyl or 1-imidazolyl group wherein said group is optionally substituted with one or two $\text{C}_1\text{-C}_4$ alkyl groups;

R^7 is H or $\text{C}_1\text{-C}_4$ alkyl;

and R^8 is H; $\text{C}_1\text{-C}_3$ alkyl or (hydroxy) $\text{C}_2\text{-C}_3$ alkyl;

a compound of formula (III):



wherein R^1 is H; $\text{C}_1\text{-C}_4$ alkyl; $\text{C}_1\text{-C}_4$ alkoxy or CONR^5R^6 ;
 R^2 is H or $\text{C}_1\text{-C}_4$ alkyl;
 R^3 is $\text{C}_2\text{-C}_4$ alkyl;
 R^4 is H; $\text{C}_2\text{-C}_4$ alkanoyl optionally substituted with NR^7R^8 ; (hydroxy) $\text{C}_2\text{-C}_4$ alkyl optionally substituted with NR^7R^8 ; $\text{CH}=\text{CHCO}_2\text{R}^9$; $\text{CH}=\text{CHCONR}^7\text{R}^8$; $\text{CH}_2\text{CH}_2\text{CO}_2\text{R}^9$; $\text{CH}_2\text{CH}_2\text{CONR}^7\text{R}^8$; $\text{SO}_2\text{NR}^7\text{R}^8$; $\text{SO}_2\text{NH}(\text{CH}_2)_n\text{NR}^7\text{R}^8$ or imidazolyl;
 R^5 and R^6 are each independently H or $\text{C}_1\text{-C}_4$ alkyl;
 R^7 and R^8 are each independently H or $\text{C}_1\text{-C}_4$

17

alkyl; or, together with the nitrogen atom to which they are attached, form a pyrrolidino, piperidino, morpholino or 4-(NR¹⁰)-1-piperazinyl group wherein any of said groups is optionally substituted with CONR⁵R⁶;


R⁹ is H or C₁-C₄ alkyl;

R¹⁰ is H; C₁-C₃ alkyl or (hydroxy)C₂-C₃ alkyl;

and n is 2, 3 or 4;

with the proviso that R⁴ is not H when R¹ is H, C₁-C₄ alkyl or C₁-C₄ alkoxy;

a compound of formula (IV):



wherein R¹ is C₁-C₄ alkyl;


R² is C₂-C₄ alkyl;

R³ is H or SO₂NR⁴R⁵;

R⁴ and R⁵ together with the nitrogen atom to which they are attached form a pyrrolidino, piperidino, morpholino or 4-(NR⁶)-1-piperazinyl group;

and R⁶ is H or C₁-C₃ alkyl;

or a compound of formula (V):



wherein R^1 is H; C_1 - C_4 alkyl; CN or $CONR^4R^5$;
 R^2 is C_2 - C_4 alkyl;
 R^3 is $SO_2NR^6R^7$; NO_2 ; NH_2 ; $NHCOR^8$; $NHSO_2R^8$ or $N(SO_2R^8)_2$;
 R^4 and R^5 are each independently selected from H and C_1 - C_4 alkyl;
 R^6 and R^7 are each independently selected from H and C_1 - C_4 alkyl optionally substituted with CO_2R^9 , OH, pyridyl, 5-isoxazolin-3-onyl, morpholino or 1-imidazolidin-2-onyl; or, together with the nitrogen atom to which they are attached, form a pyrrolidino, piperidino, morpholino, 1-pyrazolyl or 4-(NR^{10})-1-piperazinyl group wherein any of said groups may optionally be substituted with one or two substituents selected from C_1 - C_4 alkyl, CO_2R^9 , NH_2 and OH;
 R^8 is C_1 - C_4 alkyl or pyridyl;
 R^9 is H or C_1 - C_4 alkyl;
and R^{10} is H; C_1 - C_4 alkyl or (hydroxy) C_2 - C_3 alkyl;

or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition containing either entity, for the manufacture of a medicament for the curative or prophylactic treatment of erectile dysfunction in a male animal, including man.

2. The use according to claim 1 wherein in a compound of formula (I) R^3 is H; methyl or ethyl; R^4 is C_1-C_6 alkyl optionally substituted with cyclohexyl or with morpholino; and R^1 and R^2 are as previously defined in claim 1; in a compound of formula (II) R^1 is n-propyl; R^2 is H or methyl; R^3 is ethyl or n-propyl; R^4 is H; ethyl substituted with $CONR^5R^6$ or CO_2R^7 ; vinyl substituted with $CONR^5R^6$ or CO_2R^7 ; acetyl substituted with NR^5R^6 ; $SO_2NR^5R^6$; $CONR^5R^6$; CO_2R^7 or bromo; R^5 and R^6 together with the nitrogen atom to which they are attached form a morpholino, 4-(NR^8)-1-piperazinyl or 2,4-dimethyl-1-imidazolyl group; R^7 is H or t-butyl; and R^8 is methyl or 2-hydroxyethyl; in a compound of formula (III) R^1 is H; methyl; methoxy or $CONR^5R^6$; R^2 is H or methyl; R^3 is ethyl or n-propyl; R^4 is H; acetyl optionally substituted with NR^7R^8 ; hydroxyethyl substituted with NR^7R^8 ; $CH=CHCO_2R^9$; $CH=CHCONR^7R^8$; $CH_2CH_2CO_2R^9$; $SO_2NR^7R^8$; $SO_2NH(CH_2)_3NR^7R^8$ or 1-imidazolyl; R^5 and R^6 are each independently H or ethyl; R^7 and R^8 together with the nitrogen atom to which they are attached form a piperidino, 4-carbamoylpiperidino, morpholino or 4-(NR^{10})-1-piperazinyl group; R^9 is H or t-butyl; and R^{10} is H; methyl or 2-hydroxyethyl; with the proviso that R^4 is not H when R^1 is H, methyl or methoxy; in a compound of formula (IV) R^1 and R^2 are each independently ethyl or n-propyl; R^4 and R^5 together with the nitrogen atom to which they are attached form a 4-(NR^6)-1-piperazinyl group; and R^3 and R^6 are as previously defined in claim 1; and in a compound of formula (V) R^1 is H; n-propyl; CN or $CONH_2$; R^2 is ethyl; R^3 is $SO_2NR^6R^7$; NO_2 ; NH_2 ; $NHCOCH(CH_3)_2$;

$\text{NHSO}_2\text{CH}(\text{CH}_3)_2$; $\text{NHSO}_2(3\text{-pyridyl})$ or $\text{N}[\text{SO}_2(3\text{-pyridyl})]_2$; R^6 is H; methyl or 2-hydroxyethyl; R^7 is methyl optionally substituted with 2-pyridyl or 5-isoxazolin-3-onyl; or ethyl 2-substituted with OH, $\text{CO}_2\text{CH}_2\text{CH}_3$, morpholino or 1-imidazolidin-2-onyl; or R^6 and R^7 together with the nitrogen atom to which they are attached form a (4- CO_2R^9)piperidino, 5-amino-3-hydroxy-1-pyrazolyl or 4-(NR^{10})-1-piperazinyl group; R^9 is H or ethyl; and R^{10} is H; methyl or 2-hydroxyethyl.

3. The use according to claim 2 wherein in a compound of formula (III) R^1 is methyl; CONH_2 or $\text{CONHCH}_2\text{CH}_3$; R^2 is H; R^3 is ethyl or n-propyl; R^4 is H; acetyl; 1-hydroxy-2-(NR^7R^8)ethyl; $\text{CH}=\text{CHCO}_2\text{C}(\text{CH}_3)_3$; $\text{CH}=\text{CHCONR}^7\text{R}^8$; $\text{SO}_2\text{NR}^7\text{R}^8$ or 1-imidazolyl; R^7 and R^8 together with the nitrogen atom to which they are attached form a 4-(NR^{10})-1-piperazinyl group; and R^{10} is methyl or 2-hydroxyethyl; with the proviso that R^4 is not H when R^1 is methyl; in a compound of formula (IV) R^1 is n-propyl; R^2 is ethyl; and R^3 is 1-piperazinylsulphonyl or 4-methyl-1-piperazinylsulphonyl; and in a compound of formula (V) R^1 is n-propyl or CN; R^2 is ethyl; R^3 is $\text{SO}_2\text{NR}^6\text{R}^7$; $\text{NHSO}_2\text{CH}(\text{CH}_3)_2$; $\text{NHSO}_2(3\text{-pyridyl})$ or $\text{N}[\text{SO}_2(3\text{-pyridyl})]_2$; R^6 is H or methyl; R^7 is methyl; or ethyl 2-substituted with $\text{CO}_2\text{CH}_2\text{CH}_3$; morpholino or 1-imidazolidin-2-onyl; or R^6 and R^7 together with the nitrogen atom to which they are attached form a (4- CO_2R^9)piperidino or 4-(NR^{10})-1-piperazinyl group; R^9 is H or ethyl; and R^{10} is H; methyl or 2-hydroxyethyl.

4. The use according to claim 2 or claim 3 wherein the compound of formula (I), (II), (III), (IV) or (V) is selected from

1-ethyl-5-[5-(n-hexylsulphamoyl)-2-n-propoxy-phenyl]-3-methyl-1,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one;

1-ethyl-5-(5-diethylsulphamoyl-2-n-propoxy-phenyl)-3-methyl-1,6-dihydro-7H-pyrazolo[4,3-d]-

pyrimidin-7-one;

5-[5-(N-cyclohexylmethyl-N-methylsulphamoyl)-2-n-propoxyphenyl]-1-ethyl-3-methyl-1,6-dihydro-7H-pyrazolo[4,3-d]pyrimidin-7-one;

6-(5-bromo-2-n-propoxyphenyl)-3-methyl-1-n-propyl-1,5-dihydro-4H-pyrazolo[3,4-d]pyrimidin-4-one;

3-methyl-6-(5-morpholinosulphonyl-2-n-propoxyphenyl)-1-n-propyl-1,5-dihydro-4H-pyrazolo[3,4-d]pyrimidin-4-one;

6-[5-(2-carboxyvinyl)-2-n-propoxyphenyl]-3-methyl-1-n-propyl-1,5-dihydro-4H-pyrazolo[3,4-d]pyrimidin-4-one;

6-[5-(2-t-butoxycarbonylvinyl)-2-n-propoxyphenyl]-3-methyl-1-n-propyl-1,5-dihydro-4H-pyrazolo[3,4-d]pyrimidin-4-one;

3-methyl-6-[5-(2-morpholinocarbonylvinyl)-2-n-propoxyphenyl]-1-n-propyl-1,5-dihydro-4H-pyrazolo[3,4-d]pyrimidin-4-one;

3-methyl-6-[5-(2-morpholinocarbonylethyl)-2-n-propoxyphenyl]-1-n-propyl-1,5-dihydro-4H-pyrazolo[3,4-d]pyrimidin-4-one;

2-{2-ethoxy-5-[4-(2-hydroxyethyl)-1-piperazinylsulphonyl]phenyl}-8-methylquinazolin-4(3H)-one;

2-{5-[4-(2-hydroxyethyl)-1-piperazinylsulphonyl]-2-n-propoxyphenyl}-8-methylquinazolin-4(3H)-one;

8-methyl-2-{5-[2-(4-methyl-1-piperazinylcarbonyl)-ethenyl]-2-n-propoxyphenyl}quinazolin-4(3H)-one;

8-carbamoyl-2-{2-ethoxy-5-[4-(2-hydroxyethyl)-1-piperazinylsulphonyl]phenyl}quinazolin-4(3H)-one;

8-ethylcarbamoyl-2-(2-n-propoxyphenyl)quinazolin-4(3H)-one;

2-[2-ethoxy-5-(4-ethoxycarbonylpiperidinosulphonyl)phenyl]-8-n-propylpyrido[3,2-d]pyrimidin-4(3H)-one;

2-[5-(4-carboxypiperidinosulphonyl)-2-ethoxyphenyl]-8-n-propylpyrido[3,2-d]pyrimidin-4(3H)-one;

2-{2-ethoxy-5-[4-(2-hydroxyethyl)-1-piperazinyl-sulphonyl]phenyl}-8-n-propylpyrido[3,2-d]pyrimidin-4(3H)-one;

and 2-{2-ethoxy-5-[(bis-3-pyridylsulphonyl)amino]-phenyl}-8-n-propylpyrido[3,2-d]pyrimidin-4(3H)-one.

5. A pharmaceutical composition for the curative or prophylactic treatment of erectile dysfunction in a male animal, including man, comprising a compound of formula (I), (II), (III), (IV) or (V) according to any one of claims 1 to 4, or a pharmaceutically acceptable salt thereof, together with a pharmaceutically acceptable diluent or carrier.

6. A method of treating a male animal, including man, to cure or prevent erectile dysfunction which comprises treating said male animal with an effective amount of a compound of formula (I), (II), (III), (IV) or (V) according to any one of claims 1 to 4, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition containing either entity.

7. The use of a compound of formula (I), (II), (III), (IV) or (V) according to any one of claims 1 to 4, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition containing either entity, for the manufacture of a medicament for the curative or prophylactic treatment of female sexual dysfunction, premature labour or dysmenorrhea.

8. A method of treating a female animal, including woman, to cure or prevent sexual dysfunction, premature labour or dysmenorrhea which comprises treating said female animal with an effective amount of a compound of formula (I), (II), (III), (IV) or (V) according to any one of claims 1 to 4, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition containing either entity.

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/505

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	WO,A,94 28902 (PFIZER LTD) 22 December 1994 see claims ---	1-6
Y	WO,A,94 00453 (PFIZER LTD) 6 January 1994 cited in the application see claims 1-7 ---	1-6
Y	WO,A,93 06104 (PFIZER LTD) 1 April 1993 cited in the application see claims 1-7 ---	1-6
Y	WO,A,93 12095 (PFIZER LTD) 24 June 1993 cited in the application see claims 1-8 ---	1-6
	--- -/-	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *&* document member of the same patent family

Date of the actual completion of the international search

24 January 1996

Date of mailing of the international search report

07.02.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+ 31-70) 340-3016

Authorized officer

Klaver, T

INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 95/04065

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO,A,93 07149 (PFIZER LTD) 15 April 1993 cited in the application see claims 1-7 ---	1-6
Y	WO,A,94 05661 (PFIZER LTD) 17 March 1994 cited in the application see claims 1-7 ---	1-6
Y	AM. J. PHYSIOL. HEART CIRC. PHYSIOL., vol. 264, no. 2, 1993 pages h419-h422, F. TRIGA-ROCHA ET AL 'Nitric oxide and cGMP: mediators of pelvic nerve-stimulated erection in dogs.' see the whole document ---	1-6
A	NEUROL URODYN., vol. 13, no. 1, 1994 pages 71-80, F. TRIGO-ROCHA ET AL. 'Intracellular mechanism of penile erection in monkeys.' ---	
A	EP,A,0 535 924 (MERCK FROSST CANADA INC) 7 April 1993 -----	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 95/04065

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9428902	22-12-94	AU-B- 6797394 NO-A- 954757	03-01-95 24-11-95
WO-A-9400453	06-01-94	CA-A- 2139109 EP-A- 0647227 FI-A- 946083 JP-T- 7504681	06-01-94 12-04-95 23-12-94 25-05-95
WO-A-9306104	01-04-93	PT-A- 100862	30-11-93
WO-A-9312095	24-06-93	CA-A- 2122360 EP-A- 0628032 FI-A- 942769 JP-T- 7502029 US-A- 5482941	24-06-93 14-12-94 10-06-94 02-03-95 09-01-96
WO-A-9307149	15-04-93	PT-A- 100915	29-10-93
WO-A-9405661	17-03-94	CA-A- 2138298 EP-A- 0656898 FI-A- 950889 JP-T- 7506838	17-03-94 14-06-95 27-02-95 27-07-95
EP-A-535924	07-04-93	JP-A- 7215934 US-A- 5273980	15-08-95 28-12-93

INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 95/04065

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 6+8 are directed to a method of treatment of (diagnostic method practised on) the human/animal body the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.